

Vaccination Report – 2 August 2022

1. Vaccine Implementation

- WHO's Emergency Use Listing(EUL) Vaccines (Last Updated 7 July 2022)

	Manufacturer	Name of Vaccine	NRA of Record	Vaccine type
1	Pfizer-BioNTech (US)	BNT162b2/COMIRNATY Tozinameran (INN)	EMA,USFDA	mRNA
2	AstraZeneca (UK)	AZD1222 Vaxzevria	EMA, MFDS KOREA, Japan MHLW/PMDA, Australia TGA, COFEPRIS(Mexico), ANMAT(Argentina)	Non ReplicatingViral vector
3	Serum Institute of India (India)	Covishield (ChAdOx1_nCoV-19)	DCGI	Non Replicating Viral Vector
4	Johnson &Johnson (US)	Ad26.CoV2.S	EMA, DCGI	Non ReplicatingViral vector
5	Moderna (US)	mRNA-1273	EMA, USFDA, MFDS	mRNA
6	Sinopharm Beijing (China)	SARS-CoV-2 Vaccine(Vero Cells)	NMPA	Inactivated virus (Vero Cells)
7	Sinovac (China)	COVID-19 Vaccine (Vero Cells)	NMPA	Inactivated virus (Vero Cell)
8	Bharat Biotech (India)	SARS-CoV-2 Vaccine, Inactivated (Vero Cell)/ COVAXIN	DCGI	Whole-Virion Inactivated (Vero Cell)
9	Serum Institute of India (India)	NVX-CoV2373/Covovax	DCGI	Protein Subunit
10	NOVAVAX (US)	NVX-CoV2373/Covovax	EMA	Protein Subunit
11	CanSinoBIO (China)	Ad5-nCoV	NMPA	Non ReplicatingViral vector

- **40** Vaccines Approved by at Least One Country

Vaccine Type	mRNA	Non Replicating Viral vector	Inactivated virus	Protein Subunit	DNA	Virus-like Particles (VLP)	Total
In Use	4	7	11	16	1	1	40

Source: <https://covid19.trackvaccines.org/vaccines/> (Last Updated 2 August, 2022)

- Vaccination against COVID-19 has now started in **218** locations

(Source: [Our World in Data](#). Last Updated 1 August, 2022)

Location	Doses Given	Complete Initial Protocol (% of population)	Partly Vaccinated (% of population)
Worldwide	12.35 billion	4.87 billion (61.59 %)	5.30 billion (67.04 %)

About this data:


a: This data changes rapidly and might not reflect doses still being reported. It may differ from other sites & sources.

b: Where data for full vaccinations is available, it shows how many people have received at least 1 dose and how many people have been fully vaccinated (which may require more than 1 dose). Where data for full vaccinations isn't available, the data shows the total number of vaccine doses given to people. Since some vaccines require more than 1 dose, the number of fully vaccinated people is likely lower.

c: It only has full vaccination totals in some locations.

Share of people vaccinated against COVID-19, Aug 1, 2022

Our World in Data



Source: Official data collated by Our World in Data


Note: Alternative definitions of a full vaccination, e.g. having been infected with SARS-CoV-2 and having 1 dose of a 2-dose protocol, are ignored to maximize comparability between countries.

CC BY

Share of people who received at least one dose of COVID-19 vaccine, Aug 1, 2022

Our World in Data

Total number of people who received at least one vaccine dose, divided by the total population of the country.




Source: Official data collated by Our World in Data – Last updated 2 August 2022

OurWorldInData.org/coronavirus • CC BY

COVID-19 vaccine boosters administered per 100 people, Aug 1, 2022

Total number of vaccine booster doses administered, divided by the total population of the country. Booster doses are doses administered beyond those prescribed by the original vaccination protocol.




Source: Official data collated by Our World in Data – Last updated 2 August 2022

OurWorldInData.org/coronavirus • CC BY

COVID-19 vaccination policy, Aug 1, 2022

Policies for vaccine delivery. Vulnerable groups include key workers, the clinically vulnerable, and the elderly. "Others" include select broad groups, such as by age.



Source: Oxford COVID-19 Government Response Tracker, Blavatnik School of Government, University of Oxford – Last updated 2 August 2022
OurWorldInData.org/coronavirus • CC BY

2. Effectiveness of Vaccine and/or Previous Infection against symptomatic infection for Alpha, Delta and Omicron variants

Vaccine Status	Vaccine Effectiveness		
	Alpha	Delta	Omicron
1 Dose (BNT162b2 or ChAdOx1 nCoV-19)	48.7% (95%CI: 45.5-51.7%) ¹ 66%(BNT162b2) ⁴ 64% (ChAdOx1) ⁴	30.7% (95%CI: 25.2-35.7%) ¹ 56%(BNT162b2) ⁴ 67%(ChAdOx1) ⁴ 82% (95% CI:73- 91%) ⁷	
1 Dose (mRNA-1273)	83% ⁴	72% ⁴	
1 Dose(Sinopharm or Sinovac)		13.8%,(95%CI: -60.2-54.8%) ³	

2 Doses (BNT162b2)	93.7% (95%CI: 91.6-95.3) ¹ 76% (95%CI: 69-81%) ² 89% ⁴	88% (95%CI: 85.3-90.1%) ¹ 42% (95% CI: 13-62%) ² 87% ⁴ 93% (95% CI: 88-97%/12-18Y) ⁵ 93% (95% CI: 88-97%) ⁷	50% (95% CI: 35%–62%) ⁸
2 Doses (ChAdOx1 nCoV-19)	74.5% (95%CI: 68.4-79.4%) ¹	67.0% (95%CI: 61.3-71.8%) ¹	
2 Doses (mRNA-1273)	86% , (95%CI: 81-90.6%) ²	76% , (95% CI: 58-87%) ²	30.4% (95% CI: 5.0%-49.0%) ⁹
2 Doses(Sinopharm or Sinovac)		59.0% , (95%CI: 16.0-81.6%) ³	
3 Doses (BNT162b2)		95.33% (SD 6.44) ⁶ 86.1% (95% CI, 67.3 to 94.1) ¹¹	67.2% (95% CI: 66.5- 67.8%) at 2 to 4 weeks ¹⁰ 49.4% (95% CI, 47.1 to 51.6) ¹¹ 52.2% (95% CI, 48.1 to 55.9) ¹²
3 Doses(mRNA-1273)			62.5% (95% CI: 56.2-67.9%) ⁹ 47.3% (95% CI, 40.7 to 53.3) ¹¹
2 Doses (BNT162b2) + 1Dose(mRNA-1273)			73.9% (95% CI: 73.1- 74.6%) at 2 to 4 weeks ¹⁰
2 Doses(ChAdOx1 nCoV-19)+1Dose(BNT162b2)			62.4% (95% CI, 61.8- 63.0) at 2 to 4 weeks ¹⁰
2 Doses (ChAdOx1 nCoV-19)+ 1Dose (mRNA-1273)			70.1% (95% CI, 69.5 to 70.7) at 2 to 4 weeks ¹⁰
2 Doses (BNT162b2) +Previous infection			55.1% (95% CI, 50.9 to 58.9) ¹²
3 Doses (BNT162b2) +Previous infection			77.3% (95% CI, 72.4 to 81.4) ¹²
Previous Omicron Infection			76.1% on BA.4 or BA.5 (95% CI: 54.9 to 87.3%) ¹³

References:

- 1) [Effectiveness of Covid-19 Vaccines against the B.1.617.2 \(Delta\) Variant](#)
- 2) [Comparison of two highly-effective mRNA vaccines for COVID-19 during periods of Alpha and Delta variant prevalence](#)
- 3) [Efficacy of inactivated SARS-CoV-2 vaccines against the Delta variant infection in Guangzhou: A test-negative case-control real-world study](#)
- 4) [Effectiveness of COVID-19 vaccines against variants of concern in Ontario, Canada](#)
- 5) [Effectiveness of BNT162b2 Vaccine against Delta Variant in Adolescents](#)
- 6) [A RCT of a third dose CoronaVac or BNT162b2 vaccine in adults with two doses of CoronaVac](#)
- 7) [Effectiveness of BNT162b2 Vaccine against Delta Variant in Adolescents](#)
- 8) [Effectiveness of BNT162b2 Vaccine against Omicron Variant in South Africa](#)
- 9) [Effectiveness of mRNA-1273 against SARS-CoV-2 omicron and delta variants](#)
- 10) [Covid-19 Vaccine Effectiveness against the Omicron \(B.1.1.529\) Variant](#)
- 11) [Effect of mRNA Vaccine Boosters against SARS-CoV-2 Omicron Infection in Qatar](#)
- 12) [Effects of Previous Infection and Vaccination on Symptomatic Omicron Infections](#)
- 13) [Protection of SARS-CoV-2 natural infection against reinfection with the BA.4 or BA.5 Omicron subvariants](#)

3. Latest Relevant Articles

- [Neutralization of SARS-CoV-2 Omicron sublineages by 4 doses of mRNA vaccine](#) (Published: 29 July 2022)

- Comparative effectiveness of BNT162b2 versus mRNA-1273 boosting in England: a cohort study in OpenSAFELY-TPP (Published: July 30, 2022)
- Antibody Response Following COVID-19 Boosters During the Omicron Wave in the United States: A Decentralized, Digital Health, Real-World Study (Published: July 31, 2022)
- Association of Receiving a Fourth Dose of the BNT162b Vaccine With SARS-CoV-2 Infection Among Health Care Workers in Israel (Published August 2, 2022)

4. Other Information

- CDC: Safety Monitoring of COVID-19 mRNA Vaccine Second Booster Doses Among Adults Aged ≥ 50 Years — United States, March 29, 2022–July 10, 2022 (July 29, 2022)